



Anti-fibrotic effect of bovine colostrum in carbon tetrachloride-induced hepatic fibrosis

Dong Hyun Sinn, Geum-Youn Gwak, Young Ju Kwon, Seung Woon Paik

Department of Medicine, Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul, Korea

Received: April 24, 2017

Revised: May 12, 2017

Accepted: May 19, 2017

Corresponding author:

Geum-Youn Gwak

Department of Medicine,
Samsung Medical Center,
Sungkyunkwan University
School of Medicine, 81 Irwon-ro,
Gangnam-gu, Seoul 06351, Korea
Tel: +82-2-3410-3409
E-mail: gy.gwak@samsung.com

ABSTRACT

Purpose: To investigate the preventive effect of colostrum on hepatic fibrosis induced by carbon tetrachloride (CCl₄) in mice.

Methods: Seven-week-old male Balb/C mice were treated with CCl₄ with or without a diet of bovine colostrum (BC) for 2 weeks. Hepatic fibrosis was determined using Sirius red staining, α -smooth muscle actin (α -SMA) immunohistochemistry, immunoblot analysis, and quantitative polymerase chain reaction (PCR) for profibrogenic and proinflammatory genes. Hepatocellular injury was assessed through the measurement of serum alanine aminotransferase (ALT) levels.

Results: Compared to the CCl₄ group, the BC+CCl₄ group had a 55% reduction in fibrosis as demonstrated by Sirius red staining. The BC+CCl₄ group also had a significant reduction in α -SMA expression according to immunohistochemistry and immunoblot. On quantitative PCR, the BC+CCl₄ group showed significant 8.42-, 21.11-, 0.57-, and 2.90-fold reductions in the expression of *Acta2*, *Col1a1*, *Tgf β 1*, and *Tnf- α* , respectively, compared to the CCl₄ group. The serum ALT level was significantly reduced in the BC+CCl₄ group compared to the CCl₄ group.

Conclusion: The present study demonstrated that BC ameliorated CCl₄-induced hepatic fibrosis by inhibiting fibrogenesis-related gene expression and also decreased CCl₄-induced hepatocyte injury. Therefore, BC might be therapeutically efficacious in preventing hepatic fibrosis in various liver diseases.

Keywords: Carbon tetrachloride; Colostrum; Fibrosis; Liver; Wounds and injuries

INTRODUCTION

Hepatic fibrosis is the result of chronic hepatocellular injury and the first step toward the development of devastating complications such as cirrhosis and hepatocellular carcinoma [1]. The process of hepatic fibrosis is considered to be reversible [2], and the blockade or reversal of hepatic fibrosis has become an important issue in medicine worldwide [3-5]. With advances in knowledge of hepatic fibrosis [6-9], considerable progress has been made in the development of novel drugs; however, most of these drugs are still under clinical trial, with uncertain efficacy and adverse events.

Colostrum is milk from lactating mammals that is secreted during the first approximately 72

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>).

hours after birth. Colostrum differs from mature milk in that it contains abundant bioactive components, including growth factors, immunoglobulins, lactoperoxidase, lysozyme, lactoferrin, vitamins, peptides, and oligosaccharides, which are of great importance to human health [10,11]. Thus, colostrum has a variety of beneficial health effects such as bacteriostatic, bactericidal, anti-viral, anti-inflammatory, and immunomodulatory protection [12,13], as well as high antioxidant capacity and radical scavenging activity [14,15].

Oxidative stress is known to be a major contributing factor to the onset of hepatic fibrosis [16]. In pathologic conditions such as chronic hepatitis B virus (HBV) infection, alcoholic liver disease, and non-alcoholic fatty liver disease, the reactive oxygen species (ROS) generated within hepatocytes up-regulate the expression of fibrosis-associated genes, which causes hepatic stellate cell (HSC) activation and collagen deposition. In an experimental setting, carbon tetrachloride (CCl₄) is able to induce hepatic fibrosis by stimulating the formation of free radicals, ROSs and lipid peroxidation products [17] and thus is widely used to generate hepatic fibrosis in model animals [18,19].

This study was designed to investigate the preventive effect of colostrum on hepatic fibrosis induced by CCl₄ in mice. The study provides a reliable experimental basis for the application of colostrum in the prevention and treatment of hepatic fibrosis.

METHODS

Animals and experimental design

Thirty male (aged 7 weeks) Balb/C mice were purchased from OrientBio (Seongnam, Korea). Following more than 7 days of acclimation, the mice were divided into four groups: the control group (n=5) was fed with water (15 μ L/g) five times a week and injected with corn oil (2 μ L/g) twice a week intraperitoneally, the bovine colostrum (BC) group (n=5) was fed with BC (15 μ L/g) five times a week and injected with corn oil (2 μ L/g) twice a week intraperitoneally, the CCl₄ group (n=10) was fed with water (15 μ L/g) five times a week and injected with CCl₄ (1:3 diluted in corn oil, 2 μ L/g) twice a week intraperitoneally, and the BC+CCl₄ group (n=10) was fed with BC (15 μ L/g) five times a week and injected with CCl₄ (1:3 diluted in corn oil, 2 μ L/g) twice a week intraperitoneally. The mice were treated for 2 weeks and sacrificed 48 hours after the final corn oil or CCl₄ injection. When the mice were sacrificed, blood samples and livers were obtained. Blood samples were collected from the inferior vena cava (IVC), centrifuged,

and stored at -80°C . All mice were given standard laboratory chow and water *ad libitum* and maintained under a 12-hour light/dark cycle. All animal experiments were performed according to the guidelines of the Center for Laboratory Animal Research of Samsung Medical Center and with the approval of the Institutional Animal Care and Use Committee of Samsung Medical Center (protocol number: 20140513001).

Materials

BC was purchased from Diotech Korea (Seoul, Korea). CCl₄ and corn oil were purchased from Sigma (St. Louis, MO, USA).

Measurements of serum alanine aminotransferase

Alanine aminotransferase (ALT) levels were measured with sera collected from the IVC using an FUJI DRI-CHEM SLIDE GPT/ALP-PIII (Fujifilm Co., Tokyo, Japan) and spectrophotometer (FUJI DRI-CHEM 3000, Fujifilm Co.), according to the manufacturer's instructions.

Histological examination and immunohistochemistry

Liver samples were fixed in 10% formalin and embedded in paraffin. The sections were then cut (5 μ m) and stained with Sirius red to assess fibrosis. For immunohistochemical analysis, the samples were incubated with monoclonal antibody to α -smooth muscle actin (α -SMA) (1:500 dilution, Sigma).

RNA isolation and quantitative polymerase chain reaction analysis

RNA was isolated with a combination of Trizol and subsequent Qiagen RNeasy Plus Mini Kit according to the manufacturer's instructions (Qiagen, Hilden, Germany). Following reverse transcription, quantitative real-time polymerase chain reaction (qPCR) was performed on an ABI 7300 cycler using TaqMan probe sets for α -actin-2 (*Acta2*), α -1 type I collagen (*Col1a1*), tissue inhibitor of metalloproteinase-1 (*Timp1*), transforming growth factor β 1 (*Tgf β 1*), interleukin 6 (*Il6*), tumor necrosis factor α (*Tnf- α*), and glyceraldehyde 3-phosphate dehydrogenase (*GAPDH*) (Applied Biosystems, Foster City, CA, USA). The expression of each gene was normalized to *GAPDH* and fold-induction was calculated based on the $\Delta\Delta\text{Ct}$ method.

Western blot

We performed electrophoresis of protein extracts using T-PER Tissue Protein Extraction Reagent (Thermo Fisher Scientific Inc., Hanover Park, IL, USA) and subsequent blotting. For Western blotting, proteins were transferred to nitrocellulose mem-



PRECISION AND FUTURE MEDICINE

Anti-fibrotic effect of bovine colostrum



branes using an iBLOT blotting system (Life Technologies, Seoul, Korea). We incubated blots with mouse antibody to α -SMA (Sigma) at a dilution of 1:500. Chemiluminescence signals were detected using SuperSignal West Pico Chemiluminescent Substrate (Thermo Fisher Scientific Inc.). Blots were probed with mouse antibody to β -actin (Sigma).

Statistical analysis


All data are expressed as mean \pm standard deviation. Differences between experimental and control groups were assessed using two-tailed unpaired Student t-tests. P-values less than 0.05 were considered statistically significant.



A



B



C

Fig. 1. Bovine colostrum (BC) ameliorates carbon tetrachloride (CCl₄)-induced hepatic fibrosis. Male Balb/c mice were injected with corn oil or CCl₄ with or without BC treatment. Hepatic fibrosis was evaluated using (A) Sirius red staining, (B) α -smooth muscle actin (α -SMA) immunohistochemistry, and (C) α -SMA immunoblotting.

RESULTS

Forty 8 hours after the final CCl₄ injection, the CCl₄ group showed overt hepatic fibrosis induction by 53% compared to the control group, whereas the BC+CCl₄ group had a 55% reduction in fibrosis as demonstrated by Sirius red staining (Fig. 1A).

In the analysis of α -SMA expression, which is a marker of activated HSCs, CCl₄ significantly increased α -SMA expression levels compared to those in the control group, whereas additional BC treatment reduced α -SMA expression to the same level as in the control group, as demonstrated by immunohistochemistry and immunoblotting (Fig. 1B, C).

We then analyzed the contribution of BC to the fibrogenic response of the liver. CCl₄ treatment increased the expression of *Acta2*, *Col1a1*, *Timp1*, and *Tgf β 1*, which encode profibrogenic markers, and of proinflammatory cytokine genes *Il6* and *Tnf- α* (Fig. 2). Compared to the CCl₄ group, the BC+CCl₄ group showed significant 8.42-, 21.11-, 0.57-, and 2.90-fold reductions in the expression of *Acta2*, *Col1a1*, *Tgf β 1*, and *Tnf-*

alpha, respectively ($P < 0.05$) (Fig. 2). The *Timp1* and *Il6* expression were reduced by 4.51- and 4.33-fold in the BC+CCl₄ group versus the CCl₄ group, respectively; however, the differences did




Fig. 3. Bovine colostrum (BC) inhibits hepatocellular injury. Hepatocellular injury was determined using serum alanine aminotransferase (ALT) level. ALT level was measured using sera collected from the inferior vena cava. CCl₄, carbon tetrachloride.




Fig. 2. Bovine colostrum (BC) inhibits fibrogenesis-related gene expression. Levels of *Acta2*, *Col1a1*, *Timp1*, *Tgf β 1*, *Il6*, and *Tnf- α* mRNA were determined using quantitative real-time polymerase chain reaction. Gene expression is expressed as fold-induction in comparison to the control group. CCl₄, carbon tetrachloride; *Acta2*, α -actin-2; *Col1a1*, α -1 type I collagen; *TIMP1*, tissue inhibitor of metalloproteinase-1; *Tgf β 1*, transforming growth factor β 1; *Il6*, interleukin 6; *Tnf- α* , tumor necrosis factor α .

not reach statistical significance.

We also found a significant reduction in hepatocellular injury as determined by serum ALT levels in the BC+CCl₄ group compared to the CCl₄ group (Fig. 3).

DISCUSSION

Oxidative stress is a major factor contributing to the onset of hepatic fibrosis in various liver diseases, including chronic HBV infection and long-term alcohol consumption [16]. ROS are generated mainly via the mitochondrial electron transport chain or via activation of cytochrome P450, mostly cytochrome P450 2E1, nicotinamide adenine dinucleotide phosphate oxidase, or xanthine oxidase or mitochondrial damage. ROS generation in HSC and myofibroblasts occurs in response to several known profibrogenic mediators, including angiotensin II, platelet-derived growth factor, TGF β , and leptin [20], and the generated ROS up-regulate the expression of critical fibrosis-related genes such as COL1A1, COL1A2, monocyte chemoattractant protein-1 (MCP1), and TIMP1 via activation of signal transduction pathways and transcription factors, including c-Jun N-terminal kinase, activator protein-1, and nuclear factor- κ B [1]. Oxidative stress also alters mitochondrial permeability, which leads to the promotion of hepatocyte necrosis and/or apoptosis. Furthermore, hepatocarcinogenesis in the cirrhotic liver is associated with severe oxidative stress [21].

In an experimental setting, CCl₄ induces hepatic fibrosis via the formation of free radicals, ROS, and lipid peroxidation products [17]. CCl₄ is metabolized into trichloromethyl (-CCl₃) and trichloromethyl peroxy (-OCCl) by cytochrome P450 enzymes in the liver, and generates free radicals in the process. These free radicals are responsible for oxidative stress-induced hepatic damage and fibrosis [22]. Thus, the CCl₄-induced fibrosis model is widely used to evaluate the hepatoprotective effects of various pharmacological therapies. Previous reports have demonstrated that several antioxidant drugs can attenuate CCl₄-induced hepatotoxicity [23,24]. The administration of N-acetylcysteine decreases liver fibrosis by increasing glutathione (GSH) level and decreasing the expression of induced nitric oxide synthase (iNOS) [23]. Moreover, treatment with insulin-like growth factor-I reduced myeloperoxidase level and the expression of iNOS in CCl₄ model rats [24]. In the present study, we investigated the preventive effect of BC on hepatic fibrosis induced by CCl₄. BC dramatically ameliorated CCl₄-induced hepatic fibrosis by inhibiting fibrogenesis-related gene expression and also decreased CCl₄-

induced hepatic injury.

To date, many studies have demonstrated the protective effect of breast milk against various infectious diseases in infants, including respiratory infections, diarrhea, otitis media, and infections in premature infants [25,26]. In addition to the numerous clinically significant benefits of breastfeeding, breast-milk also has bioactive components that protect newborns from hyperoxic challenge due to transition of life to an environment far richer in oxygen than the intrauterine environment [27,28]. It is believed that oxidative stress is involved in the pathogenesis of numerous neonatal diseases such as bronchopulmonary dysplasia, retinopathy of prematurity and necrotizing enterocolitis [29,30]. Breast milk has a number of enzymatic and non-enzymatic antioxidant constituents, including superoxide dismutase, GSH peroxidase, catalase, vitamin E, vitamin C, and β -carotene, which protect newborns against ROS in the early stage of life [29,31]. Accordingly, previous studies have demonstrated the high antioxidant capacity and radical scavenging activity of colostrum. Zarban et al. [14] measured total antioxidant capacity (TAC) and free radical scavenging activity in human milk during the first 6 months of lactation. Both TAC and free radical scavenging activity were significantly higher in colostrum than transitional and mature milk. Quiles et al. [15] also showed that the TAC of milk was significantly higher in colostrum compared with mature and transitional milk. Thus, colostrum might be therapeutically useful for various pathologic conditions in which oxidative stress is involved in the pathogenesis.

In summary, the present study demonstrated that colostrum can ameliorate CCl₄-induced hepatic fibrosis by inhibiting fibrogenesis-related gene expression and also decreases CCl₄-induced hepatic injury. Therefore, colostrum might be therapeutically efficacious in preventing hepatic fibrosis in a variety of human liver diseases.

CONFLICTS OF INTEREST

No potential conflict of interest relevant to this article was reported.

ACKNOWLEDGMENTS

This research was supported by the Basic Science Research Program through the National Research Foundation of Korea (NRF) funded by the Ministry of Science, ICT & Future Planning (grant number: 2012R1A1A3010961).

ORCID

Geum-Youn Gwak <https://orcid.org/0000-0002-6453-3450>

REFERENCES

- Bataller R, Brenner DA. Liver fibrosis. *J Clin Invest* 2005; 115:209-18.
- Povero D, Busletta C, Novo E, di Bonzo LV, Cannito S, Paternostro C, et al. Liver fibrosis: a dynamic and potentially reversible process. *Histol Histopathol* 2010;25:1075-91.
- Svegliati Baroni G, D'Ambrosio L, Ferretti G, Casini A, Di Sario A, Salzano R, et al. Fibrogenic effect of oxidative stress on rat hepatic stellate cells. *Hepatology* 1998;27:720-6.
- Hellerbrand C, Stefanovic B, Giordano F, Burchardt ER, Brenner DA. The role of TGFbeta1 in initiating hepatic stellate cell activation in vivo. *J Hepatol* 1999;30:77-87.
- Tsukamoto H. Is interleukin-10 antifibrogenic in chronic liver injury? *Hepatology* 1998;28:1707-9.
- Knittel T, Janneck T, Muller L, Fellmer P, Ramadori G. Transforming growth factor beta 1-regulated gene expression of Ito cells. *Hepatology* 1996;24:352-60.
- Pinzani M, Milani S, Grappone C, Weber FL Jr, Gentilini P, Abboud HE. Expression of platelet-derived growth factor in a model of acute liver injury. *Hepatology* 1994;19:701-7.
- Lasky JA, Ortiz LA, Tonthat B, Hoyle GW, Corti M, Athas G, et al. Connective tissue growth factor mRNA expression is upregulated in bleomycin-induced lung fibrosis. *Am J Physiol* 1998;275(2 Pt 1):L365-71.
- Abraham DJ, Shiwen X, Black CM, Sa S, Xu Y, Leask A. Tumor necrosis factor alpha suppresses the induction of connective tissue growth factor by transforming growth factor-beta in normal and scleroderma fibroblasts. *J Biol Chem* 2000;275:15220-5.
- Mero A, Miiikulainen H, Riski J, Pakkanen R, Aalto J, Takala T. Effects of bovine colostrum supplementation on serum IGF-I, IgG, hormone, and saliva IgA during training. *J Appl Physiol* (1985) 1997;83:1144-51.
- Severin S, Wenshui X. Milk biologically active components as nutraceuticals: review. *Crit Rev Food Sci Nutr* 2005;45: 645-56.
- Buescher ES, McWilliams-Koeppen P. Soluble tumor necrosis factor-alpha (TNF-alpha) receptors in human colostrum and milk bind to TNF-alpha and neutralize TNF-alpha bioactivity. *Pediatr Res* 1998;44:37-42.
- Garofalo RP, Goldman AS. Cytokines, chemokines, and colony-stimulating factors in human milk: the 1997 update. *Biol Neonate* 1998;74:134-42.
- Zarban A, Taheri F, Chahkandi T, Sharifzadeh G, Khorshadizadeh M. Antioxidant and radical scavenging activity of human colostrum, transitional and mature milk. *J Clin Biochem Nutr* 2009;45:150-4.
- Quiles JL, Ochoa JJ, Ramirez-Tortosa MC, Linde J, Bompadre S, Battino M, et al. Coenzyme Q concentration and total antioxidant capacity of human milk at different stages of lactation in mothers of preterm and full-term infants. *Free Radic Res* 2006;40:199-206.
- Muriel P. Role of free radicals in liver diseases. *Hepatol Int* 2009;3:526-36.
- Weber LW, Boll M, Stampfl A. Hepatotoxicity and mechanism of action of haloalkanes: carbon tetrachloride as a toxicological model. *Crit Rev Toxicol* 2003;33:105-36.
- Pierce RA, Glaug MR, Greco RS, Mackenzie JW, Boyd CD, Deak SB. Increased procollagen mRNA levels in carbon tetrachloride-induced liver fibrosis in rats. *J Biol Chem* 1987;262:1652-8.
- Hernandez-Munoz R, Diaz-Munoz M, Suarez J, Chagoya de Sanchez V. Adenosine partially prevents cirrhosis induced by carbon tetrachloride in rats. *Hepatology* 1990; 12:242-8.
- De Minicis S, Bataller R, Brenner DA. NADPH oxidase in the liver: defensive, offensive, or fibrogenic? *Gastroenterology* 2006;131:272-5.
- Shimamoto K, Hayashi H, Taniai E, Morita R, Imaoka M, Ishii Y, et al. Antioxidant N-acetyl-L-cysteine (NAC) supplementation reduces reactive oxygen species (ROS)-mediated hepatocellular tumor promotion of indole-3-carbinol (I3C) in rats. *J Toxicol Sci* 2011;36:775-86.
- Jimenez W, Claria J, Arroyo V, Rodes J. Carbon tetrachloride induced cirrhosis in rats: a useful tool for investigating the pathogenesis of ascites in chronic liver disease. *J Gastroenterol Hepatol* 1992;7:90-7.
- Pereira-Filho G, Ferreira C, Schwengber A, Marroni C, Zettler C, Marroni N. Role of N-acetylcysteine on fibrosis and oxidative stress in cirrhotic rats. *Arq Gastroenterol* 2008;45: 156-62.
- Garcia-Fernandez M, Castilla-Cortazar I, Diaz-Sanchez M, Navarro I, Puche JE, Castilla A, et al. Antioxidant effects of insulin-like growth factor-I (IGF-I) in rats with advanced liver cirrhosis. *BMC Gastroenterol* 2005;5:7.
- Lonnerdal B. Breast milk: a truly functional food. *Nutrition* 2000;16:509-11.
- Lawrence RM, Pane CA. Human breast milk: current concepts of immunology and infectious diseases. *Curr Probl*

PRECISION AND FUTURE MEDICINE

Anti-fibrotic effect of bovine colostrum

- Pediatr Adolesc Health Care 2007;37:7-36.
27. Saugstad OD. Bronchopulmonary dysplasia-oxidative stress and antioxidants. *Semin Neonatol* 2003;8:39-49.
 28. Friel JK, Friesen RW, Harding SV, Roberts LJ. Evidence of oxidative stress in full-term healthy infants. *Pediatr Res* 2004;56:878-82.
 29. Scheibmeir HD, Christensen K, Whitaker SH, Jegaethesan J, Clancy R, Pierce JD. A review of free radicals and antioxidants for critical care nurses. *Intensive Crit Care Nurs* 2005;21:24-8.
 30. Thompson AM, Bizzarro MJ. Necrotizing enterocolitis in newborns: pathogenesis, prevention and management. *Drugs* 2008;68:1227-38.
 31. L'Abbe MR, Friel JK. Superoxide dismutase and glutathione peroxidase content of human milk from mothers of premature and full-term infants during the first 3 months of lactation. *J Pediatr Gastroenterol Nutr* 2000;31:270-4.