



Advances in biomarker for stroke patients: from marker to regulator

Oh Young Bang^{1,2}

¹Department of Neurology, Samsung Medical Center, Sungkyunkwan University School of Medicine, Seoul, Korea

²Translational and Stem Cell Research Laboratory on Stroke, Samsung Medical Center, Seoul, Korea

Received: January 16, 2017

Revised: February 26, 2017

Accepted: February 27, 2017

Corresponding author:

Oh Young Bang

Department of Neurology,
Samsung Medical Center,
Sungkyunkwan University
School of Medicine, 81 Irwon-ro,
Gangnam-gu, Seoul 06351, Korea
Tel: +82-2-3410-3599
E-mail: ohyoung.bang@samsung.com

ABSTRACT

Biomarkers refer to indicators found in the blood, other body fluids or tissues that predict physiologic or disease states, increased disease risk, or pharmacologic responses to a therapeutic intervention. Stroke is a heterogeneous condition, and stroke biomarkers could be used as a guiding tool for more effective personalized therapy. In this review, the recent advances in the biomarkers in stroke field are discussed. First, various types of biomarkers including genetic, extracellular vesicle, and metabolomics-associated biomarkers as well as protein biomarkers were recently introduced. The studies reviewed herein suggest that comprehensive analysis of different types of stroke biomarkers will improve the understanding of individual pathophysiologies and further promote the development of screening tool of high-risk patients, predicting model of stroke outcome and rational stroke therapy tailored to the characteristics of each case. Second, several biomarkers can be bio-‘makers’ that regulate compensatory or pathological process in the development of stroke etiology and recovery after stroke. Several protein (e.g., chemokines, caveoli), genetic (e.g., microRNA), and extracellular vesicles (e.g., cancer cell, stem cells-derived) may be directly involved in these processes. These bio-makers may be molecular target of treatment and can be used for new drug development.

Keywords: Biomarkers; Precision medicine; Risk factors; Stroke, ischemic; Therapeutics

INTRODUCTION

Precision medicine relies on biomarkers with which to better classify patients with their disease risk, prognosis, and treatment response. In a narrow sense, biomarkers refer to indicators measured by chemical or biologic tests using blood or urine that predicts physiologic or disease states, or increased disease risk. Biomarkers are also a valuable tool in drug development providing more accurate and complete information regarding drug performance, disease progression, or response to a specific drug therapy. Treatment according to the biomarkers has also been investigated in various diseases including ischemic heart disease, cancer, or immunological disorders. On the contrary, there has been relatively lack of biomarker researches in cerebrovascular disease.

The importance of considering heterogeneity among stroke patients has emerged. Unlike

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>).

coronary heart disease, stroke has heterogeneous pathophysiologies and mechanisms. Moreover, individual patients with stroke have different features even among subjects with same stroke mechanisms. These aspects enhance the need for development of personalized medicine based on characteristics of each patient rather than performing large randomized clinical trials.

Herein, we review the current and new stroke biomarkers with their strengths and weaknesses focusing on the importance of comprehensive approaches and the role of biomarkers on the development of new drugs for stroke.

ROLE OF BIOMARKERS IN STROKE RESEARCH

Screening high-risk subjects

Although many attempts including national publicity and various programs for health promotion have been made to manage stroke risk factors, the prevalence of stroke has not been markedly reduced. This may be partially resulted from hidden risk factors of stroke. Interestingly, certain regions in the United States (stroke belt and buckle) have an unusually high incidence and mortality of stroke and the phenomenon could not be explained by the differences of the conventional risk factors [1,2]. Therefore, many researchers have devoted themselves to find nontraditional risk factors of stroke to explain it and numerous possible contributing factors have been identified, including metabolic syndrome, sleep-related breathing disorders, and air pollution [3,4].

In addition to these nontraditional risk factors, a series of biomarkers reflecting inflammation, hemostasis, thrombosis, endothelial function, or neurohormonal activity have been evaluated as potential tools in an effort to improve risk prediction of future stroke, and thereby avert future events [5-12]. Our recent biomarker study showed that although traditional risk factors have been reported to be different between cerebral microangiopathy and macroangiopathy, endothelial dysfunction and related renal dysfunction were associated with both types of cerebral angiopathies [13]. In the near future, genome-wide association study may also greatly contribute to build risk stratification models by identifying genetic variants that confer susceptibility to cerebrovascular disease [14].

Rapid stroke diagnosis

Although the diagnosis of acute stroke mostly relies on neuroimaging techniques, the evaluation of biomarkers of tissue

injury would be an alternative strategy for rapid stroke assessment, especially for pre-hospital screening, and fast-tracking in emergency room. A rapid diagnosis of stroke based on biomarkers may be useful especially for pre-hospital screening, facilitating entry into fast track care pathway, and ancillary data when contemplating thrombolysis. However, a widely available, rapid, and sensitive diagnostic test for acute cerebral ischemia has not been available until now.

Recently, a biomarker panel rather than a single marker in isolation has been increasingly used to improve the diagnostic accuracy of suspected stroke. For instance, a diagnostic panel incorporating the levels of matrix metalloproteinase 9 (MMP-9), B-type natriuretic peptide, D-dimer, and S100 β into a composite score enhanced sensitivity of early noncontrast computed tomography (CT) alone for acute stroke, although the diagnostic accuracy was clearly imperfect [15]. Furthermore, the approach was feasible as a point-of-care test in the emergency setting [15]. As the number of presumed biomarkers for stroke expands at an exponential rate, it would be expected to develop improved biomarker combinations for more accurate diagnosis of stroke.

Detecting of possible stroke etiology and mechanisms of ischemic injury

Several studies have focused on the use of biomarkers for detecting of possible stroke mechanisms. In addition, the molecular markers related to neuronal death can provide the information about the presence of tissue at risk of infarction [16,17].

Predicting drug response and outcome

It has been well known that different patients respond in different fashions to the same medication. Among many factors that influence the effects of drugs, it is estimated that genetic factor can account for 20% to 95% of variability in drug disposition and effects [18]. For example, previous studies revealed that *CYP2C9* and *VKORC1* genetic variants are associated with warfarin dose requirement and clinical outcomes [19,20].

Besides pharmacogenetics, several biomarkers are also contributing to the predicting drug response in patients with stroke, particularly when thrombolysis is administered. Specifically, elevated S100 β and MMP-9 which were reported as serum markers of blood-brain barrier dysfunction before thrombolysis could predict hemorrhagic transformation after thrombolysis [21-23]; whereas, baseline levels of α_2 -antiplasmin were predictive of recanalization in patients treated

Table 1. Emerging roles of stroke biomarkers

Role	Example of biomarkers
Screening high-risk (asymptomatic) subjects	CRP, fibrinogen, and inflammatory cytokines, vWF, BNP, and UACR for stroke risk [5-8,10-12] ADMA for silent brain infarcts on MRI [9] Genome-wide association studies of stroke [14]
Rapid stroke diagnosis	Protein biomarkers associated with glial and neuronal cells (S100 β , GFAP, NSE, MBP), inflammation (CRP, MMP-9, VCAM, TNF- α , IL-6, VEGF), thrombosis (vWF, D-dimer), and others (BNP, homocysteine) [15,26-28]
Detection of stroke mechanisms and molecular-targeted treatment	Mechanisms of neuronal death, e.g. glutamate, GABA [16,17] Etiologic diagnosis of ischemic stroke, e.g., atherosclerotic (inflammatory markers) vs. cardioembolic (free fatty acid or vWF) vs. cancer-coagulopathy (D-dimer) [29-33]
Predicting drug response	Pharmacogenomics for the resistance to antiplatelet agents, warfarin, statin, or thrombolysis α_2 -Antiplasmin for recanalization after thrombolysis [24] MMP-9 and S100 β for a high risk of hemorrhagic transformation after thrombolysis [21-23] S100 β for delayed infarct expansion [34,35]
Predicting outcome	Inflammatory markers for early neurological worsening [36] and poor outcome after stroke [37] D-dimer for early recurrent ischemic lesion [38] Fibrin-monomer for thrombus formation in the left atrial appendage [39] vWF for adverse events or vascular events in A-fib patients receiving warfarin [40] or aspirin [41] BNP for functional outcome after A-fib stroke [42] BDNF polymorphism for outcome after traumatic brain injury or subarachnoid hemorrhage [43,44] Endothelial EV for predicting future cardiovascular events
Use as surrogate endpoints in clinical trials	Troponin T, CRP, BNP [45] S100 β , nitric oxide for phase III trials
Use as new drug development (theranostics)	Chemokines (such as SDF-1 α) to improve recovery after stroke [25] Caveolin to improve angiogenesis MicroRNA to improve recovery after stroke Cancer cell-derived EV for coagulopathy and stem cells-derived EV for stroke recovery

CRP, C-reactive protein; vWF, von Willebrand factor; BNP, B-type natriuretic peptide; UACR, urinary albumin/creatinine ratio; ADMA, asymmetrical dimethylarginine; MRI, magnetic resonance imaging; GFAP, glial fibrillary acidic protein; NSE, neuron-specific enolase; MBP, myelin basic protein; MMP-9, matrix metalloproteinase-9; VCAM, vascular cell adhesion molecule; TNF- α , tumor necrosis factor- α ; IL-6, interleukin 6; VEGF, vascular endothelial growth factor; GABA, γ -aminobutyric acid; A-fib, atrial fibrillation; BDNF, brain-derived neurotrophic factor; EV, extracellular vesicle; SDF-1 α , stromal cell-derived factor 1 α .

with thrombolysis [24]. There have been accumulating evidences that a number of biomarkers can predict clinical or radiological outcomes from cerebral ischemic events (Table 1) [5-12,14-17,21-45].

TYPE OF BIOMARKERS IN STROKE RESEARCH

Stroke biomarkers include protein, genetic, extracellular vesicle (EV), and metabolomics-associated biomarkers. Each biomarker has different aspects, and its own advantages and drawbacks (Table 2). A recent study reported that multiple EV biomarkers in addition to existing protein biomarkers are valuable for predicting future cardio- and cerebrovascular events [46]. Therefore, comprehensive approach using a variety of biomarkers is warranted to overcome the limitations. In addition, multidisciplinary approaches including neuroim-

aging biomarkers are needed.

Protein biomarkers

Researches using protein biomarkers in patients with ischemic cerebrovascular disease have mainly focused on pathophysiology, diagnosis, prognostication, and neuronal death in stroke [47]. A typical example of protein biomarkers is C-reactive protein [7,48,49]. However, a recent study raised the possibility that the relation may result from various biases [50]. Moreover, it has become skeptical about efficacy of biomarkers to predict stroke risk because they provide only limited additional information compared to the well-known stroke risk factors [51,52]. Further studies with more systematic approach and analysis are needed in this area.

Genetic biomarkers

Many epidemiological studies suggested that stroke has ge-

Table 2. Strengths and weaknesses of various biomarkers

Type	Advantage	Disadvantage
Protein	Post-stroke analysis Gene-environment interaction Quantitation	Time-dependent and need for serial tests Small numbers of final products (not causative in most)
Gene	Pre-stroke analysis Automatism Plentiful candidate Possible role of pharmacogenomics on drug effectiveness and toxicity	Influence of environment (>gene) on the development of most disease (polygenic) Clinical usefulness of genetic risk factor is low Stroke subtype-specific
MicroRNAs	Both pre- and post-stroke analysis Stably expressed in circulation for several months after stroke onset. Functional and causative	One microRNA has multiple gene targets, and need to examine the combinatorial effect of multiple microRNAs The sample size was usually limited, studies with larger patient cohorts are needed
Extracellular vesicle	Both pre- and post-stroke analysis Functional (it contains nucleic components as well as trophic factors and receptors) May be causative Window for not only damaged but also intact cells and tissues	Methods for measurement are unsettled Effortful Limited and non-specific markers for identification of extracellular vesicles
Metabolomics	Relatively plentiful candidate Monitoring of treatment effects May be functional	Effortful and assay not readily available Difficult to interpret the results (absence of clear cutoff of disease/normal state and intuitive connection between identified species and disease state) Studies in stroke field is relatively lacked

netic susceptibility, and various genetic factors were investigated [53]. However, genome-wide association studies failed to reproduce the positive results obtained from previous studies [54] or the clinical usefulness was very low [55]. For example, the hazard ratio and population attributable risk of hypertension to ischemic stroke is 2.0 and 26%, respectively. Conversely, the genetic influence on stroke was only 1.3 to 1.33 and 11% to 12%, respectively [55]. Genetic risk factors seem to be subtype-sensitive, and differential genetic risk factors have been reported to atherosclerotic, cardioembolic, and lacunar stroke [56]. Further stroke genetic studies might result in better risk prediction for different stroke subtypes and recurrent events, although conflicting result exists [57].

Recently, studies in pharmacogenomic area have been actively carried out. Among them, aspirin, clopidogrel, warfarin, statin, and thrombolytics-related genetic polymorphisms are particularly of interest. It is expected that selecting the type or dose of medication or avoiding side effects or drug resistance may be guided by simple genetic tests in the near future.

Metabolomics

The assumption of metabolomics is that occurrence of the

disease is directly related to the specific change of biochemical composition in the cell or biological fluid. Metabolomics-associated biomarker research analyzes profiles of fatty acid, amino acid, or polyamine in the blood or urine and determines normal or pathologic states. Furthermore, metabolomics-associated biomarkers can be applied to the monitoring recovery after treatment. Unfortunately, studies using metabolomics in the area of stroke is relatively lacked.

THERANOSTICS AND BIO-MAKER FOR THE DEVELOPMENT OF NEW DRUGS FOR STROKE

Theranostic nanoparticles that simultaneously deliver both imaging and therapeutic agents have gained significant attention for disease management in recent years. Theranostic agents are able to simultaneously deliver imaging agent and therapeutic drugs. Recent advances in nanomedicine offers new tool for theranostics, several studies using nanoparticles have been applied in stroke animal models. For example, ceria nanoparticles and HSP72-vectorized immunoliposome to protect against ischemic stroke, and fibrin-targeted gold nanoparticles to direct imaging of cerebral thromboemboli

on CT scan [58-60]. However, no theranostic drug have been approved by U.S. Food and Drug Administration for safety concerns.

Innate bioactive biomarker molecules may have pharmacological functions (“bio-maker”) and can be used in stroke patients. While nanomedicine-based theranostics provides artificial membrane/compounds, these natural components may role as “safe tools” for theranostics. Therefore, identification and utilization of biomarkers is vitally important for the successful development of disease modifying drugs for diseases. There have been efforts for screening candidates for new treatment strategy among biomarkers to provide a tool for new drug development for stroke. Biochemical products arising from biosynthesis of materials could protect against ischemic injury and recovery after stroke (not from breakdown of brain tissue). Several proteins (e.g., chemokines), genetic (e.g., microRNA), and EVs could be the candidates for bio-maker that regulate compensatory or pathological process in stroke. In addition, several biomarkers (e.g., serum albumin) are related to stroke outcomes and may have a role

as natural antioxidants in human. Table 3 shows essential features and examples for natural bio-makers in stroke field.

Natural antioxidants

Albumin is the most prominent protein in serum and a naturally existing antioxidant. In rat model, high-dose albumin has protective effects by reduction of the volumes of brain infarction/swelling and improvement of local perfusion to zones of critical blood flow reduction [61]. Beside albumin, bilirubin, and uric acid have been suggested to have a role as natural antioxidants in the body. Decreased levels of albumin and bilirubin were associated with poor outcome and exogenous application of albumin and uric acid may be beneficial in a certain subgroup of stroke patients (Table 4) [62-70].

MicroRNAs

MicroRNAs are short non-coding RNAs (18–23 nucleotide). MicroRNAs regress target gene expression by mRNA degradation and translation inhibition. Dysregulation of microRNAs has been linked to variety of disease development. Since many

Table 3. Essential features and examples of bio-makers for stroke

Aspect	Essential feature	Example
Roles	Diagnosis (biomarker)+Therapeutics (pharmacologic effects)	Hypoalbuminemia predicts further stroke, and application of albumin reduces infarct size
Source	Naturally exist in human body	Protein (e.g., albumin, chemokines) RNAs (e.g., microRNAs) EVs (e.g., stem cell-derived)
Molecules	Increase by compensatory/rescue mechanisms Decrease by pathophysiologic mechanism	MicroRNA related to recovery/protection Antioxidants and proteins related pathophysiology
Feasibility	Can be regulated/replaced safely with administration	Proteins or EVs

EV, extracellular vesicle.

Table 4. Natural antioxidant in the body

	Albumin	Bilirubin	Uric acid
Blood levels	Related to renal loss of albumin	Reflect intensity of oxidative stress	Decrease with oxidative stress (consumptive?) [62]
Low level	Poor outcome	Poor outcome [63] Increased incidence [64]	Controversial
High level	Not associated	Decreased incidence [65]	Not associated
Clinical trials of exogenous application	Success in a small trial (especially in recanalized patients) [66] but failed in a multicenter RCT (ALIAS trial) [67]	Not done	Failed in a RCT (URICO-ICTUS trial) [68] but improved outcome in selected patients (women or recanalized patients) [69,70]

RCT, randomized clinical trial; ALIAS, albumin treatment for acute ischaemic stroke; URICO-ICTUS, efficacy study of combined treatment with uric acid and r-tPA in acute ischemic stroke.

cell types express a set of microRNAs, microRNA signatures in circulation appear to be tissue-specific, representing their original source. Detectable levels of microRNAs exist in serum with tremendous stability, and circulating microRNAs in the serum can provide a novel class of biomarkers in cancer and other diseases [71]. In stroke field, increased or decreased levels of several circulating microRNA were reported in ischemic stroke patients [72,73]. For example, microRNA-424 level was decreased in acute stroke patients and in an animal model of stroke, and intracerebroventricular injection of lentiviral microRNA-424 reduced ischemic brain injury [72]. On the contrary, post-stroke increase in microRNA-200c contributed to ischemic brain injury by inhibiting reelin expression and inhibition of microRNA-200c resulted in an increase in cell survival [74]. There have been efforts of inhibition and overexpression of microRNAs to attenuate pathologic responses in cardiovascular disease, such as microRNA modulation using antagoMiRs (synthetic reverse compliments of oligonucleotides) that bind and silence target microRNAs or by using microRNA mimicry/pre-microRNAs that perform similarly to endogenous microRNAs [75]. Therefore, microRNAs serve as biomarker for stroke diagnosis and outcomes and also as stroke therapies.

Extracellular vesicles

EVs are defined as a heterogeneous population of small vesicles with a diameter of 0.1 to 1 μm . EVs may be a window for target cell/organs, and include genetic information (i.e., microRNAs) as well as protein inside them [76]. Moreover, it has been identified that EVs have their own function, revealing that EVs from ischemic tissue facilitated vasculogenesis in the ischemic limb model [77]. In this regard, biomarker research using EVs is a prominent field.

Nevertheless, biomarker studies using EVs in stroke are mostly performed in small cohorts. As the methods for analyzing EVs are complicated and not unique mainly due to their very small size, investigations with EVs are currently at a rudimentary state of development. When seeing the results from a large-scale clinical study for prediction of future risk of myocardial infarction, EVs could be a good candidate to compensate limitations of existing biomarker researches [46].

EVs play a critical role in the exchange of information between cells (Fig. 1). For example, stem cell-derived EVs alter the behavior of the target (damaged) cells. In recent studies, EVs secreted from mesenchymal stem cells (MSCs) promoted sciatic nerve regeneration in rats [78]. On the contrary, astrocytes release exosomes enriched in heat shock proteins and synapsin I under stress conditions [79]. Along with others, we




Fig. 1. Diagram outlining inter-cellular communications of extracellular vesicles (EVs) between neurogenic niche and damage cells. EVs released from neural progenitor/stem cells (NSCs) to extracellular milieu promote neuroprotection and neurorestoration after stroke. EVs contain genetic materials, protein and lipid and migration of EVs in the blood vessel and cerebrospinal fluid (CSF) provide the window for tissue in stroke patients.

have shown that intravenous administration of EVs derived from MSC culture media promotes functional recovery and neurovascular plasticity after stroke in rats [80,81]. No studies have examined the effects of stem cell-derived EVs in stroke patients, but a phase 1 study of cord blood-derived MSC EVs in diabetes patients is ongoing (Clinical trial identifier NCT-02138331).

Others

Caveolin

Caveolae are 50- to 100-nm cell surface plasma membrane invaginations that are abundant in endothelial cells and play a major role in the regulation of endothelial vesicular trafficking and signal transduction [82]. Our recent study showed that the serum level of caveolin-1 is decreased in patients with moyamoya disease [83]. Caveolin could be a candidate of a stroke therapy as well as a stroke biomarker. Caveolin-1 is reportedly involved in the pathogenesis of cancers and vascular diseases [82]. Caveolin-1 expression was critical for vascular endothelial growth factor-induced angiogenesis [84] and recruitment of endothelial progenitor cells from the bone marrow [85]. Caveolin expression could be modulated by genetic regulation targeting caveolin-1 using antisense/small interfering RNA or microRNA, anti-caveolin-1 antibodies, and viral vectors or polymer that target the caveolae.

Chemokines

Our previous results have shown that serum level of chemokine (stromal cell-derived factor 1α [SDF-1α]) influences the effects of neurorestorative therapies in stroke patients [86]. The levels of chemokines increase markedly in the infarcted brain during the acute phase of stroke, but decrease over time. Increase the level of SDF-1α in infarcted brain, i.e., target delivery using polymer, could modify the microenvironment to increase innate neurorestorative processes [25].

CONCLUSION

A number of biomarkers are under investigation in patients with ischemic stroke. Currently, however, application of biomarkers is only recommended for research purpose. Monitoring traditional risk factors or vessel status is more efficacious than measuring biomarkers in clinical practice. Considering advantages and disadvantages of each biomarker is important for future study, and comprehensive approach using multiple biomarkers is needed.

The role of biomarker is changing (Fig. 2). It is strongly expected that the biomarkers give us a turning point for investigating pathophysiology and therapeutic mechanisms of ischemic stroke. Continuous efforts are needed to find new biomarker and screening candidates for innate bio-makers




Fig. 2. Overview of biomarkers of stroke. (A) Changes in the role of biomarkers in stroke, (B) biomarker, bio-maker, and theranostics.

among these biomarkers is important for the development of new therapeutic strategy in stroke

CONFLICTS OF INTEREST

No potential conflict of interest relevant to this article was reported.

ORCID

Oh Young Bang <http://orcid.org/0000-0002-7962-8751>

REFERENCES

- Casper ML, Wing S, Anda RF, Knowles M, Pollard RA. The shifting stroke belt. Changes in the geographic pattern of stroke mortality in the United States, 1962 to 1988. *Stroke* 1995;26:755-60.
- El-Saed A, Kuller LH, Newman AB, Lopez O, Costantino J, McTigue K, et al. Geographic variations in stroke incidence and mortality among older populations in four US communities. *Stroke* 2006;37:1975-9.
- Howard G, Prineas R, Moy C, Cushman M, Kellum M, Temple E, et al. Racial and geographic differences in awareness, treatment, and control of hypertension: the REasons for Geographic And Racial Differences in Stroke study. *Stroke* 2006;37:1171-8.
- Bang OY, Ovbiagele B, Kim JS. Nontraditional risk factors for ischemic stroke: an update. *Stroke* 2015;46:3571-8.
- Ridker PM, Brown NJ, Vaughan DE, Harrison DG, Mehta JL. Established and emerging plasma biomarkers in the prediction of first atherothrombotic events. *Circulation* 2004;109(25 Suppl 1):IV6-19.
- Woodward M, Lowe GD, Campbell DJ, Colman S, Rumley A, Chalmers J, et al. Associations of inflammatory and hemostatic variables with the risk of recurrent stroke. *Stroke* 2005;36:2143-7.
- Ridker PM, Danielson E, Fonseca FA, Genest J, Gotto AM Jr, Kastelein JJ, et al. Rosuvastatin to prevent vascular events in men and women with elevated C-reactive protein. *N Engl J Med* 2008;359:2195-207.
- Welsh P, Lowe GD, Chalmers J, Campbell DJ, Rumley A, Neal BC, et al. Associations of proinflammatory cytokines with the risk of recurrent stroke. *Stroke* 2008;39:2226-30.
- Pikula A, Boger RH, Beiser AS, Maas R, DeCarli C, Schwedhelm E, et al. Association of plasma ADMA levels with MRI markers of vascular brain injury: Framingham offspring study. *Stroke* 2009;40:2959-64.
- Park CS, Ihm SH, Yoo KD, Kim DB, Lee JM, Kim HY, et al. Relation between C-reactive protein, homocysteine levels, fibrinogen, and lipoprotein levels and leukocyte and platelet counts, and 10-year risk for cardiovascular disease among healthy adults in the USA. *Am J Cardiol* 2010;105:1284-8.
- Wieberdink RG, van Schie MC, Koudstaal PJ, Hofman A, Witteman JC, de Maat MP, et al. High von Willebrand factor levels increase the risk of stroke: the Rotterdam study. *Stroke* 2010;41:2151-6.
- Pikula A, Beiser AS, DeCarli C, Himali JJ, Debette S, Au R, et al. Multiple biomarkers and risk of clinical and subclinical vascular brain injury: the Framingham Offspring Study. *Circulation* 2012;125:2100-7.
- Bang OY, Chung JW, Ryoo S, Moon GJ, Kim GM, Chung CS, et al. Brain microangiopathy and macroangiopathy share common risk factors and biomarkers. *Atherosclerosis* 2016;246:71-7.
- Yamada Y, Fuku N, Tanaka M, Aoyagi Y, Sawabe M, Metoki N, et al. Identification of CELSR1 as a susceptibility gene for ischemic stroke in Japanese individuals by a genome-wide association study. *Atherosclerosis* 2009;207:144-9.
- Laskowitz DT, Kasner SE, Saver J, Rempel KS, Jauch EC; BRAIN Study Group. Clinical usefulness of a biomarker-based diagnostic test for acute stroke: the Biomarker Rapid Assessment in Ischemic Injury (BRAIN) study. *Stroke* 2009;40:77-85.
- Rodriguez-Yanez M, Sobrino T, Arias S, Vazquez-Herrero F, Brea D, Blanco M, et al. Early biomarkers of clinical-diffusion mismatch in acute ischemic stroke. *Stroke* 2011;42:2813-8.
- Kernagis DN, Laskowitz DT. Evolving role of biomarkers in acute cerebrovascular disease. *Ann Neurol* 2012;71:289-303.
- Kalow W, Tang BK, Endrenyi L. Hypothesis: comparisons of inter- and intra-individual variations can substitute for twin studies in drug research. *Pharmacogenetics* 1998;8:283-9.
- Higashi MK, Veenstra DL, Kondo LM, Wittkowsky AK, Srinouanprachanh SL, Farin FM, et al. Association between CYP2C9 genetic variants and anticoagulation-related outcomes during warfarin therapy. *JAMA* 2002;287:1690-8.
- Sconce EA, Khan TI, Wynne HA, Avery P, Monkhouse L, King BP, et al. The impact of CYP2C9 and VKORC1 genetic polymorphism and patient characteristics upon warfarin dose requirements: proposal for a new dosing regimen.

- Blood 2005;106:2329-33.
21. Heo JH, Kim SH, Lee KY, Kim EH, Chu CK, Nam JM. Increase in plasma matrix metalloproteinase-9 in acute stroke patients with thrombolysis failure. *Stroke* 2003;34:e48-50.
 22. Montaner J, Molina CA, Monasterio J, Abilleira S, Arenillas JF, Ribo M, et al. Matrix metalloproteinase-9 pretreatment level predicts intracranial hemorrhagic complications after thrombolysis in human stroke. *Circulation* 2003;107:598-603.
 23. Foerch C, Wunderlich MT, Dvorak F, Humpich M, Kahles T, Goertler M, et al. Elevated serum S100B levels indicate a higher risk of hemorrhagic transformation after thrombolytic therapy in acute stroke. *Stroke* 2007;38:2491-5.
 24. Marti-Fabregas J, Borrell M, Cocho D, Belvis R, Castellanos M, Montaner J, et al. Hemostatic markers of recanalization in patients with ischemic stroke treated with rt-PA. *Neurology* 2005;65:366-70.
 25. Kim DH, Seo YK, Thambi T, Moon GJ, Son JP, Li G, et al. Enhancing neurogenesis and angiogenesis with target delivery of stromal cell derived factor-1alpha using a dual ionic pH-sensitive copolymer. *Biomaterials* 2015;61:115-25.
 26. Reynolds MA, Kirchick HJ, Dahlen JR, Anderberg JM, McPherson PH, Nakamura KK, et al. Early biomarkers of stroke. *Clin Chem* 2003;49:1733-9.
 27. Jauch EC, Lindsell C, Broderick J, Fagan SC, Tilley BC, Levine SR, et al. Association of serial biochemical markers with acute ischemic stroke: the National Institute of Neurological Disorders and Stroke recombinant tissue plasminogen activator Stroke Study. *Stroke* 2006;37:2508-13.
 28. Montaner J, Rovira A, Molina CA, Arenillas JF, Ribo M, Chacon P, et al. Plasmatic level of neuroinflammatory markers predict the extent of diffusion-weighted image lesions in hyperacute stroke. *J Cereb Blood Flow Metab* 2003;23:1403-7.
 29. Bang OY, Lee PH, Yoon SR, Lee MA, Joo IS, Huh K. Inflammatory markers, rather than conventional risk factors, are different between carotid and MCA atherosclerosis. *J Neurol Neurosurg Psychiatry* 2005;76:1128-34.
 30. Iwamoto T, Kubo H, Takasaki M. Platelet activation in the cerebral circulation in different subtypes of ischemic stroke and Binswanger's disease. *Stroke* 1995;26:52-6.
 31. Hanson E, Jood K, Karlsson S, Nilsson S, Blomstrand C, Jern C. Plasma levels of von Willebrand factor in the etiologic subtypes of ischemic stroke. *J Thromb Haemost* 2011;9:275-81.
 32. Choi JY, Jung JM, Kwon DY, Park MH, Kim JH, Oh K, et al. Free fatty acid as an outcome predictor of atrial fibrillation-associated stroke. *Ann Neurol* 2016;79:317-25.
 33. Seok JM, Kim SG, Kim JW, Chung CS, Kim GM, Lee KH, et al. Coagulopathy and embolic signal in cancer patients with ischemic stroke. *Ann Neurol* 2010;68:213-9.
 34. Matsui T, Mori T, Tateishi N, Kagamiishi Y, Satoh S, Katsube N, et al. Astrocytic activation and delayed infarct expansion after permanent focal ischemia in rats. Part I: enhanced astrocytic synthesis of s-100beta in the periinfarct area precedes delayed infarct expansion. *J Cereb Blood Flow Metab* 2002;22:711-22.
 35. Foerch C, du Mesnil de Rochemont R, Singer O, Neumann-Haefelin T, Buchkremer M, Zanella FE, et al. S100B as a surrogate marker for successful clot lysis in hyperacute middle cerebral artery occlusion. *J Neurol Neurosurg Psychiatry* 2003;74:322-5.
 36. Vila N, Castillo J, Davalos A, Chamorro A. Proinflammatory cytokines and early neurological worsening in ischemic stroke. *Stroke* 2000;31:2325-9.
 37. Whiteley W, Jackson C, Lewis S, Lowe G, Rumley A, Sandercocock P, et al. Inflammatory markers and poor outcome after stroke: a prospective cohort study and systematic review of interleukin-6. *PLoS Med* 2009;6:e1000145.
 38. Kang DW, Yoo SH, Chun S, Kwon KY, Kwon SU, Koh JY, et al. Inflammatory and hemostatic biomarkers associated with early recurrent ischemic lesions in acute ischemic stroke. *Stroke* 2009;40:1653-8.
 39. Motoki H, Tomita T, Aizawa K, Kasai H, Izawa A, Kumazaki S, et al. Coagulation activity is increased in the left atria of patients with paroxysmal atrial fibrillation during the non-paroxysmal period. Comparison with chronic atrial fibrillation. *Circ J* 2009;73:1403-7.
 40. Roldan V, Marin F, Muina B, Torregrosa JM, Hernandez-Romero D, Valdes M, et al. Plasma von Willebrand factor levels are an independent risk factor for adverse events including mortality and major bleeding in anticoagulated atrial fibrillation patients. *J Am Coll Cardiol* 2011;57:2496-504.
 41. Conway DS, Pearce LA, Chin BS, Hart RG, Lip GY. Prognostic value of plasma von Willebrand factor and soluble P-selectin as indices of endothelial damage and platelet activation in 994 patients with nonvalvular atrial fibrillation. *Circulation* 2003;107:3141-5.
 42. Rost NS, Biffi A, Cloonan L, Chorba J, Kelly P, Greer D, et al. Brain natriuretic peptide predicts functional outcome in ischemic stroke. *Stroke* 2012;43:441-5.
 43. Cheeran B, Talelli P, Mori F, Koch G, Suppa A, Edwards M,

- et al. A common polymorphism in the brain-derived neurotrophic factor gene (BDNF) modulates human cortical plasticity and the response to rTMS. *J Physiol* 2008;586: 5717-25.
44. Siironen J, Juvela S, Kanarek K, Vilkki J, Hernesniemi J, Lappalainen J. The Met allele of the BDNF Val66Met polymorphism predicts poor outcome among survivors of aneurysmal subarachnoid hemorrhage. *Stroke* 2007;38: 2858-60.
45. Halim SA, Newby LK, Ohman EM. Biomarkers in cardiovascular clinical trials: past, present, future. *Clin Chem* 2012;58:45-53.
46. Nozaki T, Sugiyama S, Koga H, Sugamura K, Ohba K, Matsuzawa Y, et al. Significance of a multiple biomarkers strategy including endothelial dysfunction to improve risk stratification for cardiovascular events in patients at high risk for coronary heart disease. *J Am Coll Cardiol* 2009;54:601-8.
47. Lynch JR, Blessing R, White WD, Grocott HP, Newman MF, Laskowitz DT. Novel diagnostic test for acute stroke. *Stroke* 2004;35:57-63.
48. Ridker PM, Cushman M, Stampfer MJ, Tracy RP, Hennekens CH. Inflammation, aspirin, and the risk of cardiovascular disease in apparently healthy men. *N Engl J Med* 1997;336:973-9.
49. Van Der Meer IM, De Maat MP, Hak AE, Kiliaan AJ, Del Sol AI, Van Der Kuip DA, et al. C-reactive protein predicts progression of atherosclerosis measured at various sites in the arterial tree: the Rotterdam Study. *Stroke* 2002;33: 2750-5.
50. Hemingway H, Philipson P, Chen R, Fitzpatrick NK, Damant J, Shipley M, et al. Evaluating the quality of research into a single prognostic biomarker: a systematic review and meta-analysis of 83 studies of C-reactive protein in stable coronary artery disease. *PLoS Med* 2010;7:e1000286.
51. Hankey GJ. Potential new risk factors for ischemic stroke: what is their potential? *Stroke* 2006;37:2181-8.
52. O'Donnell MJ, Xavier D, Liu L, Zhang H, Chin SL, Rao-Melacini P, et al. Risk factors for ischaemic and intracerebral haemorrhagic stroke in 22 countries (the INTERSTROKE study): a case-control study. *Lancet* 2010;376:112-23.
53. Dichgans M. Genetics of ischaemic stroke. *Lancet Neurol* 2007;6:149-61.
54. Lanktree MB, Dichgans M, Hegele RA. Advances in genomic analysis of stroke: what have we learned and where are we headed? *Stroke* 2010;41:825-32.
55. Ikram MA, Seshadri S, Bis JC, Fornage M, DeStefano AL, Aulchenko YS, et al. Genomewide association studies of stroke. *N Engl J Med* 2009;360:1718-28.
56. Traylor M, Farrall M, Holliday EG, Sudlow C, Hopewell JC, Cheng YC, et al. Genetic risk factors for ischaemic stroke and its subtypes (the METASTROKE collaboration): a meta-analysis of genome-wide association studies. *Lancet Neurol* 2012;11:951-62.
57. Achterberg S, Kappelle LJ, de Bakker PI, Traylor M, Algra A; SMART Study Group and the METASTROKE Consortium, et al. No additional prognostic value of genetic information in the prediction of vascular events after cerebral ischemia of arterial origin: the PROMISE study. *PLoS One* 2015;10:e0119203.
58. Kim CK, Kim T, Choi IY, Soh M, Kim D, Kim YJ, et al. Ceria nanoparticles that can protect against ischemic stroke. *Angew Chem Int Ed Engl* 2012;51:11039-43.
59. Agulla J, Brea D, Campos F, Sobrino T, Argibay B, Al-Soufi W, et al. In vivo theranostics at the peri-infarct region in cerebral ischemia. *Theranostics* 2013;4:90-105.
60. Kim JY, Ryu JH, Schellingerhout D, Sun IC, Lee SK, Jeon S, et al. Direct imaging of cerebral thromboemboli using computed tomography and fibrin-targeted gold nanoparticles. *Theranostics* 2015;5:1098-114.
61. Belayev L, Saul I, Huh PW, Finotti N, Zhao W, Busto R, et al. Neuroprotective effect of high-dose albumin therapy against global ischemic brain injury in rats. *Brain Res* 1999; 845:107-11.
62. Hong JM, Bang OY, Chung CS, Joo IS, Gwag BJ, Ovbiagele B. Influence of recanalization on uric acid patterns in acute ischemic stroke. *Cerebrovasc Dis* 2010;29:431-9.
63. Pineda S, Bang OY, Saver JL, Starkman S, Yun SW, Liebeskind DS, et al. Association of serum bilirubin with ischemic stroke outcomes. *J Stroke Cerebrovasc Dis* 2008;17:147-52.
64. Kimm H, Yun JE, Jo J, Jee SH. Low serum bilirubin level as an independent predictor of stroke incidence: a prospective study in Korean men and women. *Stroke* 2009; 40:3422-7.
65. Perlstein TS, Pande RL, Creager MA, Weuve J, Beckman JA. Serum total bilirubin level, prevalent stroke, and stroke outcomes: NHANES 1999-2004. *Am J Med* 2008;121:781-8.e1.
66. Shin DH, Moon GJ, Bang OY. Albumin therapy in acute stroke patients. *J Neurol* 2007;254:870-8.
67. Ginsberg MD, Palesch YY, Hill MD, Martin RH, Moy CS, Barsan WG, et al. High-dose albumin treatment for acute ischaemic stroke (ALIAS) Part 2: a randomised, double-blind, phase 3, placebo-controlled trial. *Lancet Neurol* 2013;12:

- 1049-58.
68. Chamorro A, Amaro S, Castellanos M, Segura T, Arenillas J, Marti-Fabregas J, et al. Safety and efficacy of uric acid in patients with acute stroke (URICO-ICTUS): a randomised, double-blind phase 2b/3 trial. *Lancet Neurol* 2014;13:453-60.
69. Llull L, Laredo C, Renu A, Perez B, Vila E, Obach V, et al. Uric acid therapy improves clinical outcome in women with acute ischemic stroke. *Stroke* 2015;46:2162-7.
70. Amaro S, Llull L, Renu A, Laredo C, Perez B, Vila E, et al. Uric acid improves glucose-driven oxidative stress in human ischemic stroke. *Ann Neurol* 2015;77:775-83.
71. Chen X, Ba Y, Ma L, Cai X, Yin Y, Wang K, et al. Characterization of microRNAs in serum: a novel class of biomarkers for diagnosis of cancer and other diseases. *Cell Res* 2008;18:997-1006.
72. Zhao H, Wang J, Gao L, Wang R, Liu X, Gao Z, et al. MiRNA-424 protects against permanent focal cerebral ischemia injury in mice involving suppressing microglia activation. *Stroke* 2013;44:1706-13.
73. Sepramaniam S, Tan JR, Tan KS, DeSilva DA, Tavintharan S, Woon FP, et al. Circulating microRNAs as biomarkers of acute stroke. *Int J Mol Sci* 2014;15:1418-32.
74. Stary CM, Xu L, Sun X, Ouyang YB, White RE, Leong J, et al. MicroRNA-200c contributes to injury from transient focal cerebral ischemia by targeting Reelin. *Stroke* 2015;46:551-6.
75. van Rooij E, Olson EN. MicroRNA therapeutics for cardiovascular disease: opportunities and obstacles. *Nat Rev Drug Discov* 2012;11:860-72.
76. Kim SJ, Moon GJ, Cho YH, Kang HY, Hyung NK, Kim D, et al. Circulating mesenchymal stem cells microparticles in patients with cerebrovascular disease. *PLoS One* 2012;7:e37036.
77. Leroyer AS, Ebrahimian TG, Cochain C, Recalde A, Blanc-Brude O, Mees B, et al. Microparticles from ischemic muscle promotes postnatal vasculogenesis. *Circulation* 2009;119:2808-17.
78. Raisi A, Azizi S, Delirezh N, Heshmatian B, Farshid AA, Amiri K. The mesenchymal stem cell-derived microvesicles enhance sciatic nerve regeneration in rat: a novel approach in peripheral nerve cell therapy. *J Trauma Acute Care Surg* 2014;76:991-7.
79. Taylor AR, Robinson MB, Gifondorwa DJ, Tytell M, Milligan CE. Regulation of heat shock protein 70 release in astrocytes: role of signaling kinases. *Dev Neurobiol* 2007;67:1815-29.
80. Xin H, Li Y, Cui Y, Yang JJ, Zhang ZG, Chopp M. Systemic administration of exosomes released from mesenchymal stromal cells promote functional recovery and neurovascular plasticity after stroke in rats. *J Cereb Blood Flow Metab* 2013;33:1711-5.
81. Moon GJ, Sung JH, Kim DH, Cho YH, Band OY. Mesenchymal stem cell-derived microvesicle therapy for stroke: neurogenic/angiogenic effects and biodistribution in a rat stroke model. Abstracts from the Fourth International Meeting of ISEV, ISEV2015, Washington, D.C., USA, 23-26 April 2015. *J Extracell Vesicles* 2015;4(Suppl 1):27783.
82. Frank PG, Woodman SE, Park DS, Lisanti MP. Caveolin, caveolae, and endothelial cell function. *Arterioscler Thromb Vasc Biol* 2003;23:1161-8.
83. Bang OY, Chung JW, Kim SJ, Oh MJ, Kim SY, Cho YH, et al. Caveolin-1, Ring finger protein 213, and endothelial function in Moyamoya disease. *Int J Stroke* 2016;11:999-1008.
84. Sonveaux P, Martinive P, DeWever J, Batova Z, Daneau G, Pelat M, et al. Caveolin-1 expression is critical for vascular endothelial growth factor-induced ischemic hindlimb collateralization and nitric oxide-mediated angiogenesis. *Circ Res* 2004;95:154-61.
85. Sbaa E, Dewever J, Martinive P, Bouzin C, Frerart F, Balligand JL, et al. Caveolin plays a central role in endothelial progenitor cell mobilization and homing in SDF-1-driven postischemic vasculogenesis. *Circ Res* 2006;98:1219-27.
86. Lee JS, Hong JM, Moon GJ, Lee PH, Ahn YH, Bang OY, et al. A long-term follow-up study of intravenous autologous mesenchymal stem cell transplantation in patients with ischemic stroke. *Stem Cells* 2010;28:1099-106.